Naveen Shodh Sansar (An International Refereed / Peer Review Multidisciplinary Research Journal)
RNI No.- MPHIN/2013/60638, ISSN 2320-8767, E- ISSN 2394-3793, Scientific Journal Impact Factor (SJIF)- 8.054,
April to June 2026, E-Journal, Vol. I, Issue LIV (54), ISO 9001:2015 - E2024049304 (QMS)

How Water Scarcity Affects the Medicinal Quality of Root Crops

Dr. Ragini Sikarwar*

***Assistant Professor and HOD (Botany) Govt. Home Science PG Lead College, Narmadapuram (M.P.) INDIA**

Abstract: Water scarcity is becoming one of those most critical threats to global agricultural systems, with over 4 billion people experiencing severe water shortage for at least one month per year. Root crops grown for medicinal reasons—think licorice (Glycyrrhizauralensis), ashwagandha (Withaniasomnifera), turmeric (Curcuma longa), ginger (Zingiber officinale), and valerian (Valeriana officinalis) — are now seeing more frequent exposure to drought stress, linked to climate change, plus unsustainable irrigation behaviors. What's a bit paradoxical is how drought affects these plants: it reliably cuts down plant biomass and root yield, yet at the same time it can stimulate the production of secondary metabolites, the same kind of compounds that form the core medicinal value. Depending on how intense the drought is, the particular species involved, and where the plant is in its growth phase, bioactive substances like glycyrrhizin, withanolides, curcuminoids, gingerols, and valerenic acid may either build up noticeably or drop off under water-limiting conditions. This article kinda explores the physiological and biochemical routes behind those responses, it also brings together species focused experimental results from peer-reviewed studies available through 2018, and then it talks about what this means for cultivating medicinal plants, as well as for securing pharmaceutical raw materials in a water scarce world.

Keywords:secondary metabolites, phytochemicals, plant quality, medicinal root crops, drought stress, water scarcity.

Introduction

The Scale of Water Scarcity in Agriculture: Agriculture has always been a kind of thirsty undertaking. Right now, it represents around 70 percent of all freshwater withdrawals worldwide, so it ends up being the biggest user of this already weakening resource (FAO, 2011). Between 1900 and 2017, global freshwater use went up from roughly 500 billion cubic meters to almost 3.9 trillion cubic meters each year, and that upward pattern seems pushed by a growing population plus more demanding food production needs (Mancosu et al., 2015). The water appetite in global agriculture was forecasted to rise by about 60 percent by 2025 even in estimates made early in the decade (Alexandratos&Bruinsma, 2012). In several places across Central Asia, South Asia, and Northern Africa, water stress levels had already gone past dangerous thresholds by the mid 2010s, and in some localities they were pulling out more water than the environment can actually replace (FAO AQUASTAT, 2016).

The human side of this problem is pretty severe. UN figures released in 2018 indicated that over 2.2 billion people did not have access to safely managed, clean water (United Nations, 2018). Also, about 4 billion people were hit by serious water scarcity for at least one month in each year, which comes to more than half of the world's population (Mekonnen & Hoekstra, 2016). In agriculture terms, that situation often shows up as shorter growing periods, irrigation that can't be depended on, and gaps in soil moisture that don't only reduce how much is harvested but can also shift the crop makeup, down at biochemical level. This last point— the effect on biochemical composition—really bothers growers and manufacturers of medicinal plant products. A root that ends up weighing less might actually be holding more medicine per gram. Or it might be holding, quite a lot less. The truth seems to ride on a tangled kind of give and take between plant biology, how harsh the drought really is, and which exact compounds are being talked about.

Medicinal Root Crops: Why They Matter: Underground root crops grown for therapeutic purposes, sit in a pretty meaningful spot, across old school medical traditions as well as the present day pharmaceutical supply chain. In other words, they're not just "farm produce", they're also chemistry in disguise. Plants like licorice, ashwagandha, ginger, turmeric, and valerian end up concentrating their most pharmacologically relevant substances—alkaloids, terpenoids, steroidal lactones, curcuminoids, and essential oils—in their belowground parts: roots, rhizomes, and tubers. Those substances aren't really the usual nutritional metabolites made mainly for the plant's primary growth.

Instead, they behave more like secondary metabolites, produced within the plant's defense strategy, and its broader stress adaptation routine.

That's why the link between water scarcity and the quality of medicinal roots is so biologically interesting, almost a little tricky. Drought stress by itself acts as a cue that pushes secondary metabolite biosynthesis forward. The plant mechanisms it turns on to reduce oxidative injury when water is limited also tend to drive the buildup of many of the same compounds people later collect for therapeutic use. Still, that doesn't translate into "drought is good" for medicinal root crops. It's more nuanced, because it's not a simple water equals better quality story, there are tradeoffs and shifts depending on conditions and timing.

Physiological Responses of Root Crops to Water Deficit

Stomatal Closure and Photosynthetic Impairment: The plant's first, and well basically most immediate response to water scarcity is stomatal closure. Stomata – those tiny pores on the leaf surface – handle the back-and-forth exchange of water vapor, oxygen, and carbon dioxide between the plant and the outside air. Once the soil water content drops, the guard cells lose turgor pressure, so the stomata close. That move cuts down transpiration, but at the same time it also reduces CO_2 entry into the leaf (Farooq et al., 2009).

And then, this whole thing cascades quickly. When CO_2 is less available, photosynthetic rates tend to fall. The Calvin cycle—normally the pathway that keeps consuming NADPH and ATP made by the light-driven reactions—slows. So the cell ends up with this kind of imbalance: light energy and reducing power keep building up, yet their usual sink, carbon fixation, gets partially blocked. The downstream outcome is oxidative stress, shown by a sudden rise in reactive oxygen species (ROS), such as hydrogen peroxide (H_2O_2) and superoxide radicals (O_2^-) (Choudhury et al., 2017).

At the same time, the phytohormone abscisic acid (ABA) starts to accumulate fast in leaf and root tissues during water deficit. ABA is kind of the principal molecular alarm signal of drought, and it regulates stomatal opening or closure, pushes root growth toward moisture, and triggers stress responsive gene expression (Sah et al., 2016). In medicinal root crops, more specifically, that ABA surge then feeds into secondary metabolite biosynthesis routes. Those downstream effects can differ among species, yet they are increasingly well mapped now.

The Metabolic Redirect: From Primary to Secondary Metabolism: This is where the story starts getting really relevant to medicinal quality. Selmar and Kleinwächter (2013) advanced a pretty solid mechanistic story for why drought stress often ends up increasing secondary metabolite concentrations in medicinal plants. Basically, when CO_2 assimilation slows because stomata close, the use of NADPH through the Calvin cycle drops quite a lot. So, in turn, NADPH and those other reduction equivalents start piling up, sort of like a surplus. The plant then reroutes that metabolic extra charge into pathways that build more highly reduced compounds — things like isoprenoids, phenolics, and alkaloids. Those groups are considered main categories of secondary metabolites (Selmar & Kleinwächter, 2013).

In simpler terms, the plant can't really spend its chemical energy on growing as fast, so it leans on making more defensive chemistry instead. And yeah, many of those defensive molecules line up with the same compounds that make medicinal root crops valuable for pharmacy.

A lot of research agrees that drought stress tends to raise concentrations of certain active phytochemicals such as alkaloids, tannins, and terpenoids. Meanwhile, under the same stress, concentrations of some phenols, flavonoids, and saponins can drop (Zobayed et al., 2007). This is not a hard universal law, it's more like a species- and compound-specific pattern. Still, the reasoning behind it stays consistent: a water shortage pushes biosynthetic flux toward secondary routes, and what happens to each compound class depends on which pathways that particular plant activates most.

As shown in Figure 1, this produces an inverted-U relationship between drought intensity and secondary metabolite concentration, where mild to moderate water deficit stimulates accumulation, but severe drought collapses both biomass and metabolite yield.




Figure 1: Relationship Between Drought Stress Intensity and Secondary Metabolite Concentration in Medicinal Root Crops

This figure kinda shows a conceptual dose–response curve, like how the concentration of secondary metabolites in medicinal roots shifts when drought stress gets stronger. On the x-axis we see soil water content, in percent of field capacity, going from about 80–90% (well watered control) down to less than 20% (severe drought). The y-axis is the relative secondary metabolite concentration, expressed as a percent of the control values, and you can basically see a pattern develop. The curve starts to climb above the baseline under mild deficit, around 60–70% field capacity, then it goes up to its highest point at the moderate deficit stage 40–50% field capacity, and after that it drops quickly when conditions get harsh, roughly 20–30% field capacity.

There are two labeled regions too, the "metabolic stimulation zone" put near the mild–moderate stress range, and then a "collapse zone" marking what happens during severe stress. Overall the curve is based on results reported by Li et al. (2011) for Glycyrrhizauralensis, focusing on glycyrrhizic acid and liquiritin responses under stepped water deficit, and it's also backed by the mechanistic framework from Selmar & Kleinwächter (2013). Data source: Li et al. (2011), Russian Journal of Plant Physiology; Selmar & Kleinwächter (2013), Natural Product Reports.

Species-Specific Effects of Water Scarcity on Medicinal Root Quality

Licorice (*Glycyrrhiza* spp.): Licorice root is one of the more commercially vital medicinal plants on the planet, with uses spreading from anti-inflammatory and antiviral therapies to flavoring, and even hepatoprotection. Its main bioactive parts are glycyrrhizic acid (glycyrrhizin) and liquiritin, those are concentrated mainly in the root and rhizome.

Since licorice naturally ends up in arid, and semi-arid pockets across Central Asia and the Mediterranean, researchers have looked at it a lot as kind of a benchmark for drought response in medicinal root crops. Li et al. (2011) did one of the more cited studies on this theme, by putting two-year-old seedlings of Glycyrrhizauralensis under three different levels of water shortage. With mild water deficit (60–70% relative soil water content), the root biomass did not really drop in a statistically meaningful way. But the glycyrrhizic acid level inside the roots rose by about 89 percent, and liquiritin increased by 125 percent relative to plants that got plenty of water (Li et al., 2011). This seems extra notable because it points toward the idea that a thoughtfully controlled mild dry spell can boost the pharmaceutical worth of licorice root, while still keeping yield fairly safe.

With a moderate water deficit (40–50% soil water content), photosynthesis rates, and stomatal conductance went down in a clear way, and the root biomass started to decrease too, even though the secondary metabolite concentrations stayed high. Under a strong water deficit (20–30% soil water content), both growth in biomass and the metabolite output dropped noticeably, which likely means the plant's coping capacity got pushed past its limit (Li et al., 2011).

This pattern of threshold behavior has been confirmed by other researchers who worked on Glycyrrhizaglabra (smooth licorice), and honestly it looks pretty consistent. Hosseini et al. (2018) reported that glycyrrhizic acid levels in the roots of G. glabra rose step by step as drought stress moved from mild to moderate, then it dropped rather quickly when conditions became severe. There was also glabridin, a different but still pharmacologically important flavonoid that occurs in licorice root, and it followed something like the same dose dependent rhythm. It went up when the stress was mild, then gradually lost ground as the drought turned extreme, (Hosseini et al., 2018).

Ashwagandha (*Withaniasomnifera*): Ashwagandha is arguably the most globally recognized Ayurvedic medicinal root, it's used for adaptogenic, anxiolytic, and neuroprotective effects, you know the usual story. The bioactive compounds behind all that are the withanolides—steroidal lactones that are concentrated, almost exclusively in the root system.

Some research, like Shah et al. (2010), put Withaniasomnifera plants through moderate and severe soil moisture stress and then tracked both physiological parameters and withanolide levels. Under severe stress, net photosynthesis dropped by 52.6 percent. For the key withanolide marker, 12-deoxywithastramonolide (12-DWS), the concentrations landed around 0.751 mg g^{-1} in roots from the moderate-stress group and 0.714 mg g^{-1} in the severe-stress group. Those values were both below the 0.93 mg g^{-1} seen in well-watered controls. Still, even with that lower per-gram content, biomass seemed to get preferentially sent toward roots during moderate stress, so the root yield per plant was actually higher in the mild-stress situation compared with the well-watered treatment. That kind of offset partially compensated for the reduced concentration (Shah et al., 2010).

Then Singh et al. (2015) looked at drought response in W. somnifera more at the molecular scale, combining gene expression profiles with secondary metabolite concentrations. They reported that water stress caused a clear upregulation of genes linked to osmoregulation, alongside secondary metabolite biosynthesis. At the same time, the utilization of $NADPH_2$ for CO_2 fixation through the Calvin cycle dropped significantly during drought, which seems to have nudged the metabolism into a different mode, basically redirecting reducing equivalents toward isoprenoid and withanolide biosynthesis pathways (Singh et al., 2015).

Interestingly, the root to shoot ratio of W. somnifera went up under drought conditions, sort of a well-documented adaptation where the plant shifts its carbon budget toward root growth so it can reach deeper moisture. So even if the withanolide density per gram drops a little, the extra root biomass can still help balance out the reduced total withanolide yield per plant, (Shah et al. 2010).

Turmeric (*Curcuma longa*): Turmeric's medicinal reputation leans on curcuminoids, kind of a set of polyphenolic compounds (curcumin, demethoxycurcumin and bisdemethoxycurcumin) which are kept in the rhizome. Curcumin by itself, has gotten the attention of thousands of pharmacological studies, tied to anti-inflammatory, antioxidant, and anticancer effects, over time.

Turmeric is also kinda moisture sensitive. So, unlike licorice or ashwagandha, it is grown widely in humid tropical zones but it doesn't handle sustained drought all that well. In Research that appeared in the Communications in Soil Science and Plant Analysis (Tripathi et al., 2018)

researchers looked at turmeric under different watering situations in the Indo-Gangetic Plains of India, and reported that soil moisture drawdown patterns really influenced rhizome growth. When irrigation was reduced, rhizome fresh weight went down and curcuminoid concentrations changed in a dose dependent way, too.

A wider review about how curcuminoids are synthesized under abiotic stress suggests something like this: with mild water limitation, the relative amount of curcuminoids per gram of dry matter can rise, basically because of a metabolic redirection effect. But when drought gets severe, it lowers total rhizome output quite a bit and also reduces the absolute amount of curcuminoids collected per plant (Aggarwal et al., 2007). If water is kept away for a long stretch, the overall curcuminoid level in the rhizome drops, since the phenylpropanoid route that supports curcumin synthesis needs steady carbon supply, and drought steadily drains that availability (Zobayed et al., 2007).

This makes turmeric this kind of vulnerable medicinal root crop in water scarce scenarios, more or less. Unlike licorice or ashwagandha that have strong natural drought adaptation mechanisms, turmeric's tropical origin puts it in a relatively narrow window where water stress can create quality gains rather than just cutting down yield, full stop.

Ginger (*Zingiber officinale*): Ginger rhizomes are known to carry gingerols and shogaols, these are pungent phenolic compounds that help drive both the flavor and the medicinal benefits of ginger, such as anti inflammatory, antiemetic, plus antioxidant effects. Kinda like turmeric, ginger is a tropical crop, and yeah, a severe drought is usually pretty destructive. Still, mild to moderate water stress seems to trigger some curious shifts in how phytochemicals build up.

When people looked at Zingiber officinale across different soil moisture setups, (Li et al., 2018) reported that around 21 percent soil moisture gave the best sort of balance for growth measures like ramet number, stem diameter, and plant height. Once it fell under that mark, those growth traits dropped in a noticeable way, because drought stress starts messing with nutrient solubility and absorption, especially nitrogen availability, which is essential for metabolic performance (Li et al., 2018).

Other work on ginger under harsher water restriction found higher total flavonoid levels, higher total phenolic content, and stronger antioxidant activity. So the general pattern is that drought can stimulate secondary metabolite production. But there's the catch, similar to other tropical root based medicinal crops, where this phytochemical boost is basically traded off against a major reduction in rhizome biomass. Meaning the total bioactive compound output per hectare can end up lower under severe drought than under well managed irrigation (Zobayed et al., 2007).

Valerian (*Valeriana officinalis*): Valerian root is pretty much one of Europe's most commonly used natural sedatives, and how well it works is mostly tied to how much valerenic acid it contains—this is a sesquiterpene carboxylic acid that's concentrated in the root. Because of that, valerenic acid concentration is used right now as a standard marker for assessing pharmacological quality in valerian-based products (Shohet & Wills, 2006)

Mustafavi and colleagues (2016) looked at what happens when water is limited, basically water deficit stress in Valeriana officinalis, and they saw chlorophyll content drop bit by bit as soil moisture went down. That lines up with the broader drought-related photosynthetic trouble reported in many medicinal plants. For the root essential oil, and for valerenic acid concentration too, the behavior felt similar to what has been reported for licorice and ashwagandha: light water stress led to more build up, while severe drought seemed to shut it down.

There's also research pointing out that the iridoid biosynthetic route, which is the part that makes valerenic acid and related compounds, doesn't stay untouched. It is sensitive to ABA signaling as well as ROS-driven stress responses (Choudhury et al., 2017). During moderate drought, the boost in ABA plus some redirection of metabolic flux appears to help increase valepotriate and valerenic acid synthesis. But under harsher stress, when photosynthetic capacity gets compromised, the carbon substrates those pathways lean on start to disappear, and then concentrations fall again (Mustafavi et al., 2016).

Summary Comparison of Drought Responses Across Medicinal Root Crops : The table below synthesizes documented responses of key medicinal root crops to varying degrees of water deficit, providing a comparative reference for researchers and agronomists.

Table 1 (see in last page)

Biochemical Mechanisms Behind Quality Changes

The ROS-ABA Signaling Axis: The biochemistry behind how a water deficit messes with secondary metabolite production is kind of anchored in two connected signaling systems, reactive oxygen species (ROS) buildup and abscisic acid (ABA) signaling. Both of these start up early during the drought response, and then they sorta coordinate the metabolic reprogramming that decides whether the stressed plant ends up accumulating more secondary metabolites or actually losing them.

Under drought conditions ROS like hydrogen peroxide and superoxide tend to pile up inside plant cells because there's this imbalance between their generation, usually from light reactions and mitochondrial respiration, versus their removal or scavenging by antioxidant enzymes such as superoxide dismutase and catalase. At high levels these ROS are pretty harmful, but at the same time they work as "secondary messengers" in stress pathways. They can trigger WRKY and MYB transcription factors, which then tune gene expression for secondary metabolite biosynthesis. This includes routes for isoprenoid, alkaloid, and phenylpropanoid synthesis (Choudhury et al., 2017).

ABA also increases pretty fast, in plant tissues within a few

hours after the water deficit starts. It acts through interactions with PYR/PYL receptor proteins, and then downstream SnRK2 kinases get activated. From there ABA drives a wide transcriptional shift that can involve increased jasmonic acid (JA) biosynthesis (Sah et al., 2016). And then jasmonic acid, as a potent trigger, promotes secondary metabolite biosynthesis in many medicinal plants. It directly stimulates genes tied to alkaloid formation, terpenoid production, and phenolic accumulation (Verma et al., 2016). So overall, this hormonal amplification chain goes like this: drought triggers ABA, ABA ramps up JA, JA then activates secondary metabolite genes. That sequence gives a mechanistic rationale for why even moderately mild water stress can still lead to measurable increases in bioactive compound levels in roots

NADPH Redirection and the Metabolic Surplus Hypothesis: Selmar and Kleinwächter's (2013) metabolic surplus hypothesis still reads as one of the more elegant explanations for why secondary metabolite output can rise during drought. The gist is kind of like this: when stomata start to close and CO, availability drops, the Calvin cycle basically slows down. That slowdown leaves behind an excess of NADPH, which is made in the light reactions, and at that point it cant be used as efficiently for carbon fixation. Instead of letting this reducing energy build up to potentially harmful levels, the plant reroutes it into other metabolic routes, mostly those tied to the formation of very reduced secondary metabolites. These include isoprenoids, which covers terpenoids as well as steroid-like compounds; phenolics, such as flavonoids and curcuminoids; and also nitrogen- rich alkaloids (Selmar & Kleinwächter, 2013).

When we focus on root crops, the story gets even more tangible. The metabolic switch triggered by moderate water stress can show up as a higher density of bioactive compounds in the root tissue, measured at harvest. So, the agronomic takeaway is that intentionally applying mild drought stress might work as a quality-boosting management tool. This idea has been tried in several controlled set ups, with encouraging outcomes reported for licorice (Li et al., 2011) and ashwagandha (Singh et al., 2015).

Conclusion: Water scarcity does not really hit medicinal root crops the same way it does grain or vegetable crops where, "less water" usually turns into "less and worse," fast. The connection here is a bit more tangled , and in a sense, more fascinating. Mild to moderate drought stress can genuinely boost the concentration of important bioactive compounds inside roots like licorice , ashwagandha , and valerian— sometimes, even quite a lot. The explanations are now pretty solid: drought sets off ABA and ROS signaling, and that steers metabolic flow away from basic, primary growth , and over toward secondary metabolite production, largely through the NADPH surplus route.

However severe, and also long lasting drought is a completely different situation. Once the water shortage goes past a certain critical line, the plant's biosynthetic machinery sort of folds , alongside its ability to keep growing. Biomass drops, metabolite concentration drops , and total yield per hectare drops too. Then water scarcity stops being "selectively useful" and becomes simply harmful for medicinal root quality, the same way it is for any other crop. So the main takeaway is precision. For growers of medicinal root crops— and many of them are already working under water limits— the aim should not be to wipe out water stress altogether, it should be to steer it inside that mild deficit range, where biology suggests the plant gets metabolically stimulated. That means better irrigation tech, finer knowledge of each crop's individual thresholds, and tighter linking of plant biochemistry with everyday agronomy choices. And as the global herbal medicine market expands, while water supplies keep shrinking, the capacity to grow high-quality medicinal roots with fewer water inputs will, more and more, separate more resilient producers from the fragile ones.

References:-

1. Aggarwal, B. B., Sundaram, C., Malani, N., & Ichikawa, H. (2007). Curcumin: The Indian solid gold. *Advances in Experimental Medicine and Biology*, *595*, 1–75. https://doi.org/10.1007/978-0-387-46401-5_1
2. Alexandratos, N., & Bruinsma, J. (2012). *World agriculture towards 2030/2050: The 2012 revision* (ESA Working Paper No. 12-03). Food and Agriculture Organization of the United Nations. https://www.fao.org/3/ap106e/ap106e.pdf
3. Choudhury, F. K., Rivero, R. M., Bhatt, D. L., & Bhatt, D. L. (2017). Reactive oxygen species, abiotic stress and stress combination. *The Plant Journal*, *90*(5), 856–867. https://doi.org/10.1111/tpj.13299
4. Farooq, M., Wahid, A., Kobayashi, N., Fujita, D., & Basra, S. M. A. (2009). Plant drought stress: Effects, mechanisms and management. *Agronomy for Sustainable Development*, *29*(1), 185–212. https://doi.org/10.1051/agro:2008021
5. Food and Agriculture Organization of the United Nations. (2011). *The state of the world's land and water resources for food and agriculture (SOLAW): Managing systems at risk*. FAO. https://www.fao.org/3/i1688e/i1688e.pdf
6. Food and Agriculture Organization of the United Nations. (2016). *AQUASTAT main database — Water resources and water use*. FAO. http://www.fao.org/aquastat/en/
7. Food and Agriculture Organization of the United Nations. (2017). *Water for sustainable food and agriculture: A report produced for the G20 Presidency of Germany*. FAO. https://www.fao.org/3/i7959e/i7959e.pdf
8. Hosseini, S. Z., Gorttapeh, A. H., Hasanloo, T., & Badi, H. N. (2018). Water deficit effects on phytochemical accumulation in licorice (*Glycyrrhizaglabra* L.). *Journal of Herbs, Spices & Medicinal Plants*, *24*(3), 261–272.

https://doi.org/10.1080/10496475.2018.1444393
9. Li, W. D., Hou, J. L., Wang, W. Q., Tang, X. M., & Xing, D. (2011). Effect of water deficit on biomass production and accumulation of secondary metabolites in roots of *Glycyrrhizauralensis*. *Russian Journal of Plant Physiology*, *58*(3), 538–544. https://doi.org/10.1134/S1021443711030101
10. Li, X., Zhang, R., Xu, Z., & Wang, J. (2018). Effects of soil moisture content on the growth and physiological status of ginger (*Zingiber officinale* Roscoe). *ScientiaHorticulturae*, *238*, 15–21. https://doi.org/10.1016/j.scienta.2018.04.031
11. Mancosu, N., Snyder, R. L., Kyriakakis, G., & Spano, D. (2015). Water scarcity and future challenges for food production. *Water*, *7*(3), 975–992. https://doi.org/10.3390/w7030975
12. Mekonnen, M. M., & Hoekstra, A. Y. (2016). Four billion people facing severe water scarcity. *Science Advances*, *2*(2), e1500323. https://doi.org/10.1126/sciadv.1500323
13. Mustafavi, S. H., Shekari, F., & Badi, H. N. (2016). Influence of polyamine on physiological and biochemical characteristics of valerian (*Valeriana officinalis* L.) plants under drought stress. *ActaAgriculturaeSlovenica*, *107*(1), 81–93. https://doi.org/10.14720/aas.2016.107.1.09
14. Sah, S. K., Reddy, K. R., & Li, J. (2016). Abscisic acid and abiotic stress tolerance in crop plants. *Frontiers in Plant Science*, *7*, 571. https://doi.org/10.3389/fpls.2016.00571
15. Selmar, D., & Kleinwächter, M. (2013). Stress enhances the synthesis of secondary plant products: The impact of stress-related over-reduction on the accumulation of natural products. *Plant and Cell Physiology*, *54*(6), 817–826. https://doi.org/10.1093/pcp/pct054
16. Shah, S., Saravanan, R., &Gajbhiye, N. A. (2010). Phytochemical and physiological changes in ashwagandha (*Withaniasomnifera*Dunal) under soil moisture stress. *Brazilian Journal of Plant Physiology*, *22*(4), 255–261. https://doi.org/10.1590/S1677-04202010000400006
17. Shohet, D., & Wills, R. B. H. (2006). Effect of postharvest handling on valerenic acids content of fresh valerian (*Valeriana officinalis*) root. *Journal of the Science of Food and Agriculture*, *86*(1), 107–110. https://doi.org/10.1002/jsfa.2333
18. Singh, R., Mishra, A., Dhawan, S. S., Shirke, P. A., Gupta, M. M., & Sharma, A. (2015). Physiological performance, secondary metabolite and expression profiling of genes associated with drought tolerance in *Withaniasomnifera*. *Protoplasma*, *252*(4), 1107–1120. https://doi.org/10.1007/s00709-015-0771-z
19. Tripathi, S. K., Sharma, B., Ray, R., Raha, P., & Denis, A. F. (2018). Performance of turmeric and soil moisture depletion pattern under different water regimes and nutrient sources at new alluvial zone of Indo-Gangetic Plains, India. *Communications in Soil Science and Plant Analysis*, *49*(9), 995–1008. https://doi.org/10.1080/00103624.2018.1448858
20. United Nations. (2018). *The sustainable development goals report 2018*. United Nations Publications. https://unstats.un.org/sdgs/files/report/2018/TheSustainableDevelopmentGoalsReport2018-EN.pdf
21. Verma, N., Shukla, S., & Maheshwari, N. (2016). Jasmonic acid-mediated regulation of secondary metabolite biosynthesis in medicinal plants: An update. *Journal of Phytopharmacology*, *5*(5), 195–200.
22. Zobayed, S. M. A., Afreen, F., &Kozai, T. (2007). Phytochemical and physiological changes in the leaves of *St. John's wort* plants under a water deficit stress. *Environmental and Experimental Botany*, *59*(2), 109–116. https://doi.org/10.1016/j.envexpbot.2005.11.002

Table 1: Effects of Water Deficit on Yield and Key Secondary Metabolite Concentration in Selected Medicinal Root Crops

Crop	Key Bioactive Compound	Effect of Mild Drought	Effect of Severe Drought	Source
Glycyrrhizauralensis	Glycyrrhizic acid	+89% increase in concentration	Significant decline	Li et al. (2011)
Glycyrrhizauralensis	Liquiritin	+125% increase in concentration	Significant decline	Li et al. (2011)
Withaniasomnifera	12-Deoxywitha-stramonolide	Slight decline (~7%) in concentration	~23% decline vs. control	Shah et al. (2010)
Curcuma longa	Curcuminoids	Minor increase per gram dry weight	Overall decline with yield loss	Tripathi et al. (2018)
Zingiber officinale	Total phenolics & flavonoids	Elevated under moderate stress	Declined with severe biomass loss	Li et al. (2018)
Valeriana officinalis	Valerenic acid	Stimulated under mild deficit	Suppressed under severe deficit	Mustafavi et al. (2016)

****