Naveen Shodh Sansar (An International Refereed / Peer Review Multidisciplinary Research Journal)
RNI No.- MPHIN/2013/60638, ISSN 2320-8767, E- ISSN 2394-3793, Scientific Journal Impact Factor (SJIF)- 8.054,
July to September 2026, E-Journal, Vol. I, Issue LV (55), ISO 9001:2015 - E2024049304 (QMS)

# Black Datura (*Datura metel*): A Brief Study of Its Leaves

**Dr. Akhilesh Chandra Verma***

***Government Naveen College, Kui-Kukdur, Dist. Kabirdham (C.G.) INDIA**

**Abstract:** *Datura metel*, commonly known as Black Datura, occupies a significant position in traditional botanical literature owing to its striking morphology, potent bioactive profile, and longstanding medicinal associations [1], [2]. Among its various organs, the leaf represents one of the most functionally important structures, serving not only as a primary photosynthetic unit but also as a major reservoir of secondary metabolites such as tropane alkaloids, phenolics, tannins, and related phytoconstituents. Despite the frequent mention of the species in ethnomedicinal discourse, focused academic treatments devoted specifically to leaf structure, composition, and applied significance remain comparatively limited. The present paper offers a concise yet critical study of the leaves of *Datura metel* through botanical, phytochemical, therapeutic, and toxicological perspectives [3], [4].

The study aims to examine the taxonomic identity and morphological characteristics of the leaves, evaluate their medicinal relevance in traditional systems, and assess the scientific concerns associated with their toxic nature. Special attention is given to the dual character of the plant: on one hand, its leaves have been historically employed in practices relating to respiratory discomfort, inflammatory conditions, dermal disorders, and analgesic applications; on the other hand, the presence of atropine, scopolamine, and hyoscyamine demands rigorous caution in handling and usage. This paradox between utility and toxicity makes the species an important subject of scholarly investigation.

Available literature indicates that the leaves possess notable pharmacological potential including antimicrobial, antioxidant, anti-inflammatory, bronchodilatory, and wound-healing tendencies under controlled conditions. However, indiscriminate use, inaccurate identification, and non-standardized dosage may result in severe physiological consequences [5]. Therefore, a balanced scientific approach is necessary.

The paper concludes that the leaves of *Datura metel* deserve continued interdisciplinary study in pharmacognosy, medicinal botany, and toxicology. Their traditional relevance may be strengthened through evidence-based validation, standardization protocols, and responsible therapeutic exploration [6].

**Keywords:** Black Datura, *Datura metel*, Leaf Morphology, Medicinal Plant, Phytochemistry, Toxicology, Ethnobotany, Solanaceae.

**Introduction -** Black Datura, botanically recognized as *Datura metel* L., is a prominent species of the family Solanaceae and is widely known for its ornamental appearance, narcotic potency, and medicinal history [1], [7]. The plant is distinguished by robust growth, broad leaves, trumpet-shaped flowers, and spiny capsules. In South Asian traditions it has long been associated with ritual, therapeutic, and folk-healing contexts. The vernacular names attached to the species across linguistic regions indicate its deep cultural familiarity. Yet, despite widespread recognition, the plant continues to evoke caution because of its toxic alkaloidal composition [8].

The term “Black Datura” generally refers to darker-stemmed or purple-tinged forms of *Datura metel*, which are frequently observed in cultivated and semi-wild habitats. Its dramatic visual form has often overshadowed the scientific importance of its vegetative organs, particularly the leaves, which constitute one of the most biologically active portions of the plant.

**Botanical Importance of *Datura metel:*** Within the genus *Datura*, *D. metel* is regarded as a species of notable taxonomic and pharmacological interest. It exhibits adaptive growth across tropical and subtropical climates and thrives in disturbed soils, roadsides, gardens, and marginal habitats. Botanically, the species demonstrates marked variability in pigmentation, floral morphology, and vegetative vigor. Such variability has encouraged comparative studies concerning chemotypes, ecological adaptability, and medicinal value [9].

From a phytobotanical standpoint, *Datura metel* is significant because it synthesizes diverse secondary metabolites, especially tropane alkaloids. These compounds contribute to the plant’s defense mechanisms and simultaneously account for its pharmacological and

toxic effects. Leaves often function as a major site of biosynthesis and accumulation, making them central to scientific investigation [10].

**Significance of Leaf Study:** The leaf is one of the most diagnostically useful and physiologically active organs of higher plants. In *Datura metel*, leaves are typically ovate to broadly elliptic, petiolate, and variably lobed or entire depending on growth conditions and genotype. Their venation pattern, surface texture, coloration, and odor provide useful characters for field identification. Beyond morphology, leaves are of considerable medicinal interest because they contain substantial concentrations of alkaloids and phenolic substances [11].

Traditional systems have used leaf preparations externally and internally for selected ailments, especially respiratory spasm, pain, inflammatory swellings, and skin conditions. Historical practices also include fumigation or smoke-based applications for bronchial distress. Modern scientific studies have further examined leaf extracts for antimicrobial and antioxidant potential. Consequently, leaf-centered research becomes essential for linking classical knowledge with contemporary evidence [12], [13].

**Purpose of the Present Study:** Although *Datura metel* has been discussed in ethnomedicinal and phytochemical literature, many studies treat the plant in a generalized manner without sufficient concentration on the leaf as a discrete unit of analysis. The present paper seeks to address that gap by presenting a systematic review of the leaves of Black Datura from botanical, medicinal, and safety-oriented perspectives.

The specific objectives are: (i) to outline the identity and structural characteristics of the leaves; (ii) to examine the major phytochemical tendencies reported in literature; (iii) to assess their traditional and emerging medicinal relevance; and (iv) to highlight toxicological concerns requiring scientific prudence. By integrating these dimensions, the study aims to present a concise but academically grounded understanding of one of the most debated leaves in medicinal botany [14].

**1. Taxonomic Position of *Datura metel* :** The scientific classification of *Datura metel* provides an essential framework for understanding its botanical identity, evolutionary relations, and medicinal relevance. As a member of the family Solanaceae, the species shares taxonomic affinity with several economically and pharmacologically significant plants. Its placement within the genus *Datura* is particularly important because the genus is distinguished by potent alkaloidal chemistry, characteristic floral architecture, and broad ethnobotanical significance. Accurate taxonomic recognition is indispensable not only for botanical scholarship but also for pharmacognostic standardization, as confusion with allied species may lead to serious toxicological consequences [1], [2].

**Table 1: Taxonomic Classification of *Datura metel***

| Rank | Classification |
|---|---|
| Kingdom | Plantae |
| Division | Magnoliophyta (Angiosperms) |
| Class | Magnoliopsida (Dicotyledons) |
| Order | Solanales |
| Family | Solanaceae |
| Genus | *Datura* |
| Species | *Datura metel* L. |

The above classification demonstrates that *Datura metel* belongs to a lineage known for chemically active metabolites and considerable medicinal utility. At the same time, many members of Solanaceae possess toxic constituents, requiring scientific caution in therapeutic interpretation [7].

**2. Botanical Description of Black Datura Leaves:** The leaves of Black Datura constitute one of the most diagnostically valuable and biologically active organs of the plant. They display notable variation according to age, environmental conditions, and cultivated form; however, certain structural traits remain sufficiently stable for identification. In medicinal botany, leaf morphology often serves as the first criterion for field recognition, collection, and preliminary authentication [14].

**Shape of Leaves:** Leaves are generally broad, ovate to elliptic-ovate, occasionally somewhat irregular in outline. Mature leaves may appear shallowly lobed or gently undulate, while younger leaves are comparatively smoother in contour. Their expanded lamina gives the plant a dense and vigorous appearance.


**Size and Structure:** The leaves are comparatively large, simple, and petiolate, with a soft but substantial laminar body. Under favorable growth conditions, they may attain considerable length and width, reflecting the species' luxuriant vegetative habit. The petiole is distinct and supports a spreading blade adapted for efficient light interception.

**Colour and Texture:** Leaf colour ranges from deep green to dark green, and in certain black-stemmed ornamental forms may appear more intense because of contrast with purple or dusky stems. The texture is usually soft, somewhat fleshy, and moderately smooth, though mature leaves may become slightly coarse with age.

**Margin, Apex and Venation:** Margins are entire to sinuately toothed or weakly lobed. The apex is commonly acute to acuminate, while the base tends to be rounded or slightly tapering. Venation is reticulate, with a prominent midrib and clearly spreading lateral veins, contributing both structural support and physiological transport efficiency.

**Odour and Surface Nature:** When crushed, leaves often emit a strong and characteristic odor associated with the plant's volatile and alkaloidal chemistry. The surface may be glabrous or faintly pubescent depending upon ecological conditions and variety. This sensory feature has historically aided traditional identification, though it cannot substitute scientific authentication [11].

**Figure -1**




The external morphology of the leaf reflects both adaptive efficiency and taxonomic utility. Its broad lamina enhances photosynthetic performance, while venation and petiole architecture support hydration and mechanical balance.

### 3. Habitat and Distribution

**Natural Habitat:** *Datura metel* commonly thrives in open, disturbed, and nutrient-responsive habitats. It is frequently encountered near roadsides, abandoned fields, garden margins, waste lands, village boundaries, and semi-cultivated landscapes. The plant shows remarkable resilience in soils affected by moderate disturbance and seasonal dryness. Such ecological plasticity contributes to its widespread persistence [9].

**Climatic Condition:** The species flourishes best in warm tropical and subtropical climates. It responds favorably to moderate sunlight, seasonal rainfall, and well-drained soils enriched with organic matter. Although comparatively hardy, prolonged frost and excessive shade restrict vigorous growth. High temperature zones with intermittent moisture generally support luxuriant foliage development.

**Distribution in India and Other Regions:** In India, *Datura metel* is distributed across plains, peri-urban landscapes, temple surroundings, agricultural edges, and domestic gardens in numerous states. It is especially familiar in northern, western, eastern, and southern regions where warm climatic conditions prevail. Beyond India, the species is reported across South and Southeast Asia, parts of Africa, tropical America, and several subtropical horticultural zones. Its spread has been facilitated both by ornamental cultivation and naturalized seed dispersal [3], [4].

The broad geographical presence of *Datura metel* explains the diversity of vernacular knowledge systems attached to the plant. Nevertheless, regional variation in morphology and phytochemical expression suggests the need for locality-specific study.

### 4. Traditional and Medicinal Importance of Leaves:

The leaves of *Datura metel* occupy a particularly significant position among the vegetative organs of the plant because they combine diagnostic botanical value with a long history of ethnomedicinal recognition. In numerous traditional systems, leaves were considered more accessible and practically utilizable than roots or seeds, as they could be harvested seasonally with relative ease. Their documented applications, however, were largely empirical in origin and varied widely according to region, healer tradition, and method of preparation. Modern interpretation therefore requires both historical sensitivity and scientific caution.

**Folk Uses of Leaves:** The leaves of *Datura metel* have occupied a notable place in several traditional healing systems of South Asia and other tropical regions. In village-based ethnomedicinal practices, warmed leaves were historically applied externally over painful joints, localized swellings, muscular stiffness, and chronic discomfort associated with seasonal ailments. In some communities, leaf paste was placed over restricted areas for temporary soothing action. Certain folk traditions also mention the cautious use of processed dried leaf material in respiratory distress; however, such practices are medically unsafe when undertaken without supervision because toxic exposure can occur rapidly. These historical references are best understood as cultural records rather than present-day therapeutic recommendations.

**Medicinal Relevance:** Phytochemical investigations indicate that the leaves contain tropane alkaloids, phenolic constituents, flavonoid-associated fractions, and other secondary metabolites. These compounds have drawn scholarly attention for possible antispasmodic, antimicrobial, anti-inflammatory, antioxidant, and neuromodulatory relevance. Preliminary pharmacological reports suggest that carefully standardized extracts may possess research value in controlled settings. Nevertheless, uncontrolled therapeutic use remains unsafe because the toxic threshold is narrow, variable, and influenced by plant maturity, ecology, and dosage inconsistency.

**External Applications:** Historically documented external applications include poultices, pastes, and oil-based preparations for rheumatic discomfort, neuralgic pain, insect irritation, glandular swelling, and certain chronic skin complaints. Such uses were generally localized rather than internal and were often accompanied by traditional restrictions regarding quantity and duration. Modern toxicological understanding demands that any external use should be approached cautiously, as dermal absorption, mucosal contact, and accidental secondary exposure may occur. Accordingly, historical utility should not be equated with routine safety.

**Table 2. Traditional Uses of *Datura metel* Leaves**

| Traditional Context | Leaf Preparation | Reported Purpose |
|---|---|---|
| Folk household remedy | Warm leaf compress | Joint pain and swelling |
| Rural topical | Leaf paste | Skin irritation / |

| | | |
|---|---|---|
| practice | | localized inflammation |
| Traditional smoke use | Dried processed leaf material | Historical respiratory use |
| Herbal oil preparations | Leaf-infused oil | Muscular discomfort |
| Ritual-ethno medicinal use | Symbolic/ceremonial use | Cultural significance |

**5. Toxic Nature and Precautions:** Despite its medicinal reputation, *Datura metel* is fundamentally recognized as a toxic plant. The leaves, though sometimes perceived as less hazardous than seeds, still contain potent pharmacologically active constituents capable of producing serious adverse reactions. For this reason, any scholarly treatment of the species remains incomplete without toxicological emphasis.

**Toxic Constituents in Leaves:** The leaves contain potent tropane alkaloids, chiefly atropine, hyoscyamine, and scopolamine. These compounds exert marked anticholinergic effects on the nervous system, cardiovascular function, secretory glands, and visual pathways. Their concentration may vary with season, maturity, habitat, plant variety, and physiological stage of growth, making casual dosage assumptions unreliable.

**Harmful Effects:** Improper exposure or ingestion may produce dryness of mouth, intense thirst, dilated pupils, blurred vision, rapid heartbeat, agitation, confusion, hallucinations, impaired coordination, urinary retention, hyperthermia, convulsions, and in severe instances coma. Sensitive individuals may respond strongly even to comparatively small quantities. Toxic reactions can therefore arise unpredictably and should always be treated as medical emergencies.

**Safe Handling Precautions:** Responsible botanical handling is essential in academic, horticultural, and field contexts. Preventive care is preferable to emergency treatment because symptom onset may be delayed yet progressive.

1. Avoid self-medication or experimental consumption.
2. Use gloves while handling fresh plant material in bulk.
3. Keep away from children and domestic animals.
4. Do not inhale smoke from leaves or seeds.
5. Wash hands and tools after botanical handling.
6. In suspected poisoning, seek immediate emergency medical care.

**Discussion:** The preceding observations reveal that the leaves of *Datura metel* cannot be interpreted through a single lens of utility or danger. They must instead be understood through an integrated botanical, ethnomedical, phytochemical, and toxicological framework.

The leaves of *Datura metel* represent a compelling intersection of botanical elegance and pharmacological seriousness. Morphologically, the broad ovate leaves are diagnostically significant for species recognition and field identification. Their venation, texture, margin variability, and dark green surface contribute to taxonomic interpretation. Chemically, they function as repositories of bioactive alkaloids and phenolic substances, explaining their long-standing medicinal reputation.

However, traditional esteem must be balanced with scientific caution. Many historical uses emerged before modern dose-standardization, toxicology, and evidence-based therapeutics. Consequently, the same leaf that was valued for pain relief or respiratory applications may also provoke severe poisoning when misidentified, overused, or crudely prepared.

From a research perspective, the species warrants continued study in phytochemistry, controlled pharmacology, toxic dose mapping, and safe derivative drug development. Standardization of extraction methods and validated analytical profiling remain especially important. Public awareness is equally essential so that ornamental familiarity does not lead to casual medicinal misuse or accidental poisoning.

**Conclusion:** The study of Black Datura leaves demonstrates how a single plant organ may embody ornamental presence, historical medicinal use, and toxicological complexity simultaneously.

*Datura metel* leaves possess clear botanical, ethnomedicinal, and phytochemical importance. Their traditional relevance is supported by the presence of physiologically active secondary metabolites, yet these same constituents confer substantial toxicity. Therefore, accurate identification, disciplined handling, and strict medical caution are indispensable.

Future investigations should prioritize standardized extraction, compound isolation, toxicity mapping, molecular characterization, and clinically validated applications. Comparative studies across ecological regions may further clarify chemical variation in leaf material. With responsible scholarship, the leaves of Black Datura may remain a subject of scientific value rather than public health risk.

**References :-**

1. U. Preissel and H. G. Preissel, *Brugmansia and Datura: Angel's Trumpets and Thorn Apples*. London: Firefly Books, 2002.
2. R. E. Schultes and A. Hofmann, *Plants of the Gods*. Rochester: Healing Arts Press, 1992.
3. W. Ruksiriwanich et al., "Wound healing effect of *Datura metel* leaf extracts," *Plants*, vol. 12, no. 13, 2023.
4. M. Prasathkumar et al., "Bioactivity of *Datura metel* leaves," *Arabian Journal of Chemistry*, vol. 15, 2022.
5. S. Bhattacharjee et al., "Mineral composition of *Datura* species," *Communications in Soil Science and Plant Analysis*, vol. 35, 2004.
6. Y. Cheng et al., "Withanolides from *Datura metel* leaves," *J. Pharm. Biomed. Anal.*, vol. 186, 2020.
7. P. S. Varier, *Indian Medicinal Plants*, vol. 2. Chennai: Orient Longman, 1994.

8. S. Rahmatullah et al., "Ethnobotanical studies on medicinal flora," *Weed Sci. Res.*, vol. 12, no. 3, 2006.
9. B. P. Pandey, *Economic Botany*. New Delhi: S. Chand & Co., 2003.
10. T. J. Tan et al., "Anti-inflammatory constituents from *Datura metel* leaves," *Fitoterapia*, vol. 142, 2020.
11. M. Anitha et al., "Phytochemical screening of *Datura metel*," *Int. J. Bioassays*, vol. 3, no. 11, 2014.
12. G. R. Liu et al., "Sesquiterpenoids from *Datura metel* leaves," *Natural Product Research*, vol. 35, 2021.
13. Z. Luo et al., "Natural products and immunoregulation," *Int. Immunopharmacol.*, vol. 110, 2022.
14. S. A. Chavhan et al., "Pharmacognostic review on *Datura*," *Int. J. Pharmacognosy Chinese Med.*, vol. 2, no. 4, 2018.
15. [15] F. C. Akharaiyi, "Antibacterial, phytochemical and antioxidant activities of *Datura metel*," *Int. J. PharmTech Res.*, vol. 3, no. 1, pp. 478–483.
16. S. Afsharypuor, A. Mostajeran, and R. Mokhtary, "Variation of scopolamine and atropine in different parts of *Datura metel* during development," *Planta Medica*, vol. 61, pp. 383–384, 1995.
17. M. Rahmatullah et al., "Medicinal plants used by traditional healers of Bangladesh," *Am.-Eurasian J. Sustain. Agric.*, vol. 4, no. 2, pp. 219–229, 2010.
18. R. J. Ko, "Clinical evaluation of suspected herbal poisoning," *Clin. Toxicol.*, vol. 37, no. 6, pp. 697–708, 1999.
19. P. C. A. Kam and S. Liew, "Traditional herbal medicine and anesthesia," *Anaesthesia*, vol. 57, pp. 1083–1089, 2002.
20. M. Yusuf, J. Begum, M. N. Hoque, and J. U. Chowdhury, *Medicinal Plants of Bangladesh*. Chittagong: BCSIR, 2009.

**********