Naveen Shodh Sansar (An International Refereed / Peer Review Multidisciplinary Research Journal)
RNI No.- MPHIN/2013/60638, ISSN 2320-8767, E- ISSN 2394-3793, Scientific Journal Impact Factor (SJIF)- 8.054,
April to June 2026, E-Journal, Vol. I, Issue LIV (54), ISO 9001:2015 - E2024049304 (QMS)

Graph Theory: From Euler's Foundations to Modern-Day Applications

Dr. Meenakshi Salviya (Rawal)*

***Assistant Professor (Mathematics) PMCoE Maharaja Bhoj Govt. P. G. College, Dhar (M.P.) INDIA**

Abstract - The study reviews modern developments along with prospective research avenues because graph theory continues its essential role in solving practical situations.

Introduction - Graph theory is a dynamic and evolving branch of mathematics that focuses on the study of relationships and connections between objects. Its origin can be traced back to simple puzzles, particularly those involving the connection of land areas by bridges. Although these early problems appeared trivial and recreational, they played a crucial role in stimulating mathematical curiosity and creativity. As noted by researchers like Norman L. Biggs, the beginning of graph theory were humble, often arising from puzzles designed to test ingenuity rather than to develop deep mathematical ideas. However, despite this simplicity, such Problems attracted the attention of mathematicians and gradually transformed into a subject rich in theoretical depth and complexity. Overtime, graph theory has developed into an exciting field with wide-ranging applications. It helps in understanding real-world networks such as brain networks, communication systems, and epidemiological structures.



The subject has expanded both theoretically and algorithmically, influencing various disciplines beyond mathematics. One of the most significant modern developments is the application of graph theory in biology. Researchers, including Martin Nowak at Harvard University, have explored how network structures affect evolutionary dynamics and the process of natural selection, originally proposed by Charles Darwin. This has led to the emergence of a new interdisciplinary field known as 'Evolutionary Graph Theory', which combines graph theory, probability and mathematical biology. Thus, from its modest beginnings in simple puzzles, graph theory has grown into a powerful and essential tool for analyzing complex systems across multiple domains.

The Origin of Graph Theory

Leonhard Euler: The Swiss mathematician Leonhard Euler (1707-1783), popularly known as the 'Analysis Incarnate', was undoubtedly a great genius who has contributed to many different, seemingly unrelated fields such as the theory of music, ship designs, complex analysis, number theory and so on. On 26 August 1735, he presented a paper titled 'solutio problematis and geometriam situs pertinentis' (on the solution to a problem in the geometry of position) at the St. Petersburg Academy. It was actually a simple looking puzzle set in the old Prussian city of Königsberg (called Kaliningrad (Russia) since 1946) and set on both sides of the Pregel river, the islands Kneiphof and Lomse. These two islands and the two mainland areas were connected to each other by seven bridges. The problem which later came to be known as the celebrated 'Königsberg Bridge problem' (KBP) was to find a route to travel from any of the four land areas (Figure 1), travel each of the seven bridges exactly once and return to the same land area. Euler used an ingenious argument to show that such a route is not possible. In this way the year 1736 became the birth year of graph theory. In graph theoretic terminologies, the graph of the Königsberg Bridge problem is a multi graph of order 4 and size 7 which is not Eulerian. It is not Eulerian because by the Euler-Hierholzer theorem, a connected graph has an Eulerian circuit if and only if every vertex has an even degree. But it is a less known fact that Euler in his paper never used the term 'Graph'. It was the British mathematician J.J. Sylvester (1814-1897) who used the

term 'Graph' for the first time in his paper titled Chemistry and Algebra, which appeared in Nature in 1878. Thus, what began as a simple and recreational puzzle, led to the foundation of graph theory, a powerful branch of mathematics that is now widely used to analyze complex networks in science, technology, and real-world.




Figure 1

Some Historical Problems in Graph Theory: Here are some important historical problems in graph theory that played a key role in its development:

(1) The Four Colour Problem: The problem was first proposed on October 23, 1852, by Francis Guthrie. While coloring a map of the countries of England, Whether four colours are sufficient to colour different countries in any map, if neighbouring countries are coloured differently. This was the origin of the much celebrated 'four-colour problem' (Four colour conjecture - 4CC). The Four colour conjecture (4CC), states: Four colours are sufficient to colour the different regions of any map drawn in the plane or on a sphere so that no two regions with a common boundary line are coloured with the same colour. Francis Guthrie's brother, Frederick, shared this observation with his professor, Augustus De Morgan. De Morgan was intrigued and wrote to the famous mathematician Sir William Rowan Hamilton about it. Interestingly, the text notes that De Morgan was born in Madurai, Tamil Nadu. In 1878, another celebrated British mathematician Arthur Cayley (1821–1895) publicised this problem in a meeting of the London Mathematical Society. A proof was given by the London-based barrister Alfred Kempe (1849–1922) in 1879, which remained as a proof for about eleven years. In 1890, a student at Oxford University, Percy Heawood (1861–1955), found an error in Kempe's argument and proved, essentially using the 'Kempe chains' argument, the "Five Colour Theorem: Every map can be coloured with five colours." However, reducing the number from five to four took more than eighty years. Many attempts to solve the four colour problem through various, different means, spanning about 120 year enriched not only the field of graph theory and in particular the theory of graph colouring but the whole of mathematics.

2. Chess Board problems: Chess Board problems occupy a central position in discrete mathematics, offering a rich interplay between combinatorial structures and algorithmic reasoning. Earliest such problem involving the movement of knights dates back to Guarini in 1512. A typical and well-known problem was, what is the minimum number of queens that can be placed on an n x n chess board 'dominating' the board, i.e., such that each square is either occupied by a queen or attacked by a queen? For the standard 8 x 8 chess board: it is known that this number is 5. But it is quite surprising that there are 4860 different ways that these five queens can be arranged to dominate the board. The Norwegian mathematician Øystein Ore (1899-1968) converted these problems to graph theoretic ones and initiated the study of 'Domination theory in graphs'. Determining domination numbers for various chessboard graphs remains an active area of research, involving combinatorial optimization and probabilistic methods. One of the most studied problems is the knight's tour, which seeks a sequence of moves that visits every vertex exactly once. In graph-theoretic terms, this corresponds to finding a Hamiltonian path (or cycle) in the knight graph. The 'Hamiltonian problem' named after the legendary Irish mathematician William Rowan Hamilton (1805-1865). In 1857 Hamilton invented the 'Icosian game' - to find such a cycle along the edges of a dodecahedron. He had to commercialise this game for his earnings.

3. Chemical graph theory : Chemical graph theory is a branch of mathematics that applies graph theory to the study of chemical compounds. In this approach, molecules are represented as graphs, where atoms are treated as vertices and chemical bonds as edges. In graph theory, a connected graph without cycle is called a tree. Many organic molecules, specially alkanes (saturated hydrocarbons) can be represented as tree structures because they contain only single bonds Methane, ethane, propane, etc. are some examples and they have the general chemical formula C_nH_{2n+2} . Since its planar representation is a connected graph with 3n+2 vertices and 3n+1 edges, it is a tree. An important result in this field is given by Arthur Cayley in 1889. It states that the number of labelled trees on n verticles n^{n-2}. This formula is very useful in counting possible molecular structures (isomers) of compounds. Even before the introduction of the term 'graph', Cayley considered 'diagrams' made of points labelled by atoms and connected by links into an assemblage. Such earlier studies have now ushered a new branch called chemical graph theory. It provides a powerful link between mathematics and chemistry. By representing molecules as graphs, it allows scientists to analyze complex chemical structures using mathematical tools, making it an essential area in both theoretical and applied research.

Spectral Graph Theory: Spectral Graph Theory is an important branch of mathematics that connects graph theory with linear algebra. It studies graph theory with linear algebra using matrices such as the adjacency matrix and the laplacian matrix. The eigen values of these matrices, known as the spectrum of the graph, provide deep insights into the structure and properties of graphs. In chemistry, especially in mathematical chemistry, graphs are used to represent molecules. Atoms are treated as vertices and

chemical bonds as edges. These descriptors help in understanding Chemical properties, reactivity, stability of molecules. One of the earliest approaches came from molecular orbital theory introduced by Erich Hückel in the 1930s. Later this idea was extended using graph theory by Ivan Gutman and others. The Wiener Index, Introduced by Harry Wiener in 1947, it is one of the oldest topological indices. It is defined as the sum of shortest path distances between all pairs of vertices in a graph. The Hosoya Index, Introduced by Haruo Hosoya in 1971, it counts the number of matchings in a graph.

Applications of Graph Theory:

(1) Biology and Medical Research: Neural networks used to model brain connectivity, epidemiology helps in studying disease spread using graph-based simulations, Genomics DNA sequencing and protein interaction networks use graph theory.

(**2) Computer Science and Technology:** Graph theory is widely used in computer science & technology to model and solve real-world problems. It helps in designing efficient computer networks for routing data, analyzing large datasets in data mining and allocating frequencies in mobile networks using graph colouring. In web design, it structures websites as interconnected graphs, while in language processing, it aids in syntax analysis through parse trees. Additionally, graph theory is applied in error detection and correction during code decoding and in optimizing electronic chip design by improving circuit connections.

(3) Physics and Chemistry: Chemistry graphs serve the purpose of modeling chemical compounds. In statistical biochemistry, certain sequences of cell samples can be eliminated to resolve conflicts between two sequences. Graph theory has a significant impact on physics & chemistry. Chart theory is employed to study molecules. The three dimensional arrangement of intricate artificial atomic systems can be quantitatively assessed by gathering statistics on graph theoretical characteristics concerning atom topology.

(4) Daily life: Graph theory helps in designing circuit connections such as series and parallel networks. It is useful for analyzing social relationships, understanding how information or rumors spread, and studying influence among individuals. It is also applied in cyber security to detect fraud and identify weak points in networks. It is used in transportation, GPS system, route optimization, and traffic management. Social media platforms also use graph theory to analyze user connections and suggest friends.

Conclusion: Graph theory is an extraordinary mathematical tool with extensive applications across various domains, including social networks, physics, biology, cyber security, computer science, and transportation. Its ability to model complex networks—ranging from communication systems to platforms like Facebook —makes it an essential tool for analyzing real-world problems. Thus graph theory has evolved from a theoretical curiosity into a powerful and indispensable framework for understanding connectivity, structure, and relationships in both natural and artificial systems.

References:-

1. Biggs, N. (1993), Algebraic Graph Theory, Second edition, Cambridge University Press.
2. Behzad M. and Chartrand G. (1971), Introduction to the Theory of Graphs.
3. John J. Watkins, (2004), Across the Board: The Mathematics of Chessboard Problems, Princeton University Press.
4. Thomas L. Saaty, Paul C. Kainen, (1977), The Four-Color Problem, Assaults and Conquest, McGraw-Hill International Book Company.
5. Vijayakumar Ambut (2022), Graph Theory: 1736-1936 and Beyond – Part I, The Mathematics Consortium.
6. Pushpa Nand Dhananjaymurthy B. V. (2023), Applications of Graph Theory in Biology and Construction, Int. J. of Math. Comput. Res. Vol. 11.
7. Kumar S. and Yadav R.B.S. (2016), Application of Graph Theory in Network Analysis, Int. J. Math. Trends Technol. Vol. 31.

✻✻✻✻✻✻✻✻✻✻✻✻